



COVID-19 Vaccine and Booster Administration Guidance for EMS Providers

Corresponds to October 23, 2021 NYS DOH Guidance
 EMS Contact: Mike McEvoy 518-423-6600



	Janssen/J&J	Moderna	Pfizer-BioNTech	Pfizer-BioNTech Pedi
Brand name	Janssen/J&J COVID-19 Vaccine	Moderna COVID-19 Vaccine	Comirnaty (COVID-19 Vaccine)	Comirnaty (COVID-19 Vaccine)
Type of Vaccine	Vector	mRNA	mRNA	mRNA
Preparation	2.5 mL multi-dose vial (5 doses)	5.5mL multi-dose vial (10 dose initial or 20 dose booster – do not puncture vial more than 20 times)	Multi-dose vial (6 dose) with purple cap - must dilute before use	Multi-dose vial (10 doses) with orange cap - must dilute before use
Dilution	n/a	n/a	Invert 10x (do not shake) before and after dilution with 1.8 mL preservative free 0.9% saline	Invert 10x (do not shake) before and after dilution with 1.3 mL preservative free 0.9% saline
Storage	Do not freeze – stored refrigerated until expiration	Stored frozen – can refrigerate for up to 30 days once thawed	Stored frozen – can refrigerate for up to 2 weeks once thawed	Stored frozen – can refrigerate for up to 10 weeks once thawed
Use after vial puncture	After puncture, stable for 6 hours refrigerated or 2 hours at room temp	After puncture, stable for 12 hours at room temp or refrigerated	After dilution, stable for 6 hours at room temp or refrigerated	After dilution, stable for 12 hours at room temp or refrigerated
Eligibility	18 years of age or older	18 years of age or older	12 years of age or older	5 to 11 years of age
First dose	0.5 mL IM (single dose)	0.5 mL IM	0.3 mL IM	0.2 mL IM
Second dose	n/a	0.5 mL IM given 28 days after 1 st	0.3 mL IM given 21 days after 1 st	0.2 mL IM given 21 days after 1 st
Third dose	Not recommended	0.5 mL IM given 28 days after 2 nd for moderate/severely immune compromised patients	0.3 mL IM given 28 days after 2 nd for moderate/severely immune compromised patients	Additional dose for moderately/severely immune compromised children not approved
Booster Eligibility	<ul style="list-style-type: none"> Anyone over 18 years of age with prior dose of any COVID-19 vaccine Boosters can be mixed/matched by patient preference. Recommended to boost with same mRNA vax when possible. Booster with mRNA vax when J&J was used initially unless patient insists on J&J booster. 			
Booster	0.5 mL IM given 2+ months after 1 st (prefer boost with mRNA vax)	0.25 mL IM given 6+ months after 2 nd	0.3 mL IM given 6+ months after 2 nd	Not approved/recommended
Expiration date check	https://vaxcheck.inj/	Scan QR code on the vial	3 months after date printed on vial	